

Package ‘swfdr’

January 27, 2023

Title Estimation of the science-wise false discovery rate and the false discovery rate conditional on covariates

Version 1.24.0

Author Jeffrey T. Leek, Leah Jager, Simina M. Boca, Tomasz Konopka

Maintainer Simina M. Boca <smb310@georgetown.edu>, Jeffrey T. Leek <jtleek@gmail.com>

Description This package allows users to estimate the science-wise false discovery rate from Jager and Leek, ``Empirical estimates suggest most published medical research is true," 2013, Biostatistics, using an EM approach due to the presence of rounding and censoring. It also allows users to estimate the false discovery rate conditional on covariates, using a regression framework, as per Boca and Leek, ``A direct approach to estimating false discovery rates conditional on covariates," 2018, PeerJ.

Depends R (>= 3.4)

Imports methods, splines, stats4, stats

License GPL (>= 3)

URL <https://github.com/leekgroup/swfdr>

BugReports <https://github.com/leekgroup/swfdr/issues>

Encoding UTF-8

LazyData true

RoxygenNote 7.1.1

Suggests dplyr, ggplot2, BiocStyle, knitr, qvalue, reshape2, rmarkdown, testthat

VignetteBuilder knitr

biocViews MultipleComparison, StatisticalMethod, Software

git_url <https://git.bioconductor.org/packages/swfdr>

git_branch RELEASE_3_16

git_last_commit f59fdb5

git_last_commit_date 2022-11-01

Date/Publication 2023-01-26

R topics documented:

| | |
|---------------------------------|---|
| BMI_GIANT_GWAS_sample | 2 |
| calculateSwfdr | 3 |
| get_number_decimals | 4 |
| journals_pVals | 5 |
| lm_pi0 | 5 |
| lm_qvalue | 7 |

| | |
|--------------|----------|
| Index | 9 |
|--------------|----------|

BMI_GIANT_GWAS_sample *Subset of SNPs from meta-analysis of BMI GWAS study.*

Description

A dataset containing 50,000 SNPs and results for their associations with BMI.

Usage

```
data(BMI_GIANT_GWAS_sample)
```

Format

A data frame with 50,000 rows and 9 variables:

SNP ID for SNP (single nucleotide polymorphism)

A1 Allele 1 for SNP

A2 Allele 2 for SNP

Freq_MAF_Hapmap Frequency of minor allele (MAF) in Hapmap project

b Estimated beta for association between SNP and BMI

se Estimated standard error (se) for association between SNP and BMI

p P-value for association between SNP and BMI

N Total sample size considered for association of SNP and BMI

Freq_MAF_Int_Hapmap Three approximately equal intervals for the Hapmap MAFs

Value

Object of class `tbl_df`, `tbl`, `data.frame`.

Source

https://www.broadinstitute.org/collaboration/giant/index.php/GIANT_consortium_data_files#GWAS_Anthropometric_2015_BMI

| | |
|----------------|---|
| calculateSwfdr | <i>Calculate the science-wise FDR (swfdr)</i> |
|----------------|---|

Description

Calculate the science-wise FDR (swfdr)

Usage

```
calculateSwfdr(
  pValues,
  truncated,
  rounded,
  pi0 = 0.5,
  alpha = 1,
  beta = 50,
  numEmIterations = 100
)
```

Arguments

| | |
|-----------------|--|
| pValues | Numerical vector of p-values |
| truncated | Vector of 0s and 1s with indices corresponding to those in pValues; 1 indicates that the p-values is truncated, 0 that it is not truncated |
| rounded | Vector of 0s and 1s with indices corresponding to those in pValues; 1 indicates that the p-values is rounded, 0 that it is not rounded |
| pi0 | Initial prior probability that a hypothesis is null (default is 0.5) |
| alpha | Initial value of parameter alpha from Beta(alpha, beta) true positive distribution (default is 1) |
| beta | Initial value of parameter beta from Beta(alpha, beta) true positive distribution (default is 50) |
| numEmIterations | The number of EM iterations (default is 100) |

Value

| | |
|-------|---|
| pi0 | Final value of prior probability - estimated from EM - that a hypothesis is null, i.e. estimated swfdr |
| alpha | Final value of parameter alpha - estimated from EM - from Beta(alpha, beta) true positive distribution |
| beta | Final value of parameter beta - estimated from EM - from Beta(alpha, beta) true positive distribution |
| z | Vector of expected values of the indicator of whether the p-value is null or not - estimated from EM - for the non-rounded p-values (values of NA represent the rounded p-values) |

| | |
|----|--|
| n0 | Expected number of rounded null p-values - estimated from EM - between certain cutpoints (0.005, 0.015, 0.025, 0.035, 0.045, 0.05) |
| n | Number of rounded p-values between certain cutpoints (0.005, 0.015, 0.025, 0.035, 0.045, 0.05) |

Examples

```
pVals <- runif(100)
tt <- rr <- rep(0, 100)
resSwfdr <- calculateSwfdr(pValues = pVals, truncated = tt, rounded = rr, numEmIterations=100)
```

```
get_number_decimals Get number of decimals (i.e. return total number of digits after decimal point) for any vector of numbers in [0,1) if number of decimals <= 6
```

Description

Get number of decimals (i.e. return total number of digits after decimal point) for any vector of numbers in [0,1) if number of decimals <= 6

Usage

```
get_number_decimals(x)
```

Arguments

x Numerical vector where all elements are in [0,1)

Value

Vector giving the number of decimals for each element in x if the number is <= 6; otherwise return 7 with a warning

Examples

```
#get_number_decimals(c(0.0006, 0.0750, 0.0420, 0.0031, 0.0001, 0.0100))
#get_number_decimals(c(6*10^-4, 7.5*10^-2, 4.2*10^-2, 3.1*10^-3, 10^-4, 10^-2))
#get_number_decimals(c(6.5*10^-4, 0.0100))
#get_number_decimals(c(6.5e-4, 0.0100))
#get_number_decimals(c(0.00065, 0.0100))
#get_number_decimals(c(10^-7, 10e-7, 10e-3))
```

| | |
|----------------|--|
| journals_pVals | <i>P-values from abstracts from articles in 5 biomedical journals (American Journal of Epidemiology, BMJ, JAMA, Lancet, New England Journal of Medicine), over 11 years (2000-2010).</i> |
|----------------|--|

Description

A dataset containing 15,653 p-values.

Usage

journals_pVals

Format

A tbl data frame with 15,653 rows and 5 variables:

pvalue P-value

pvalueTruncated Equals to 1 if the p-value is truncated, 0 otherwise

pubmedID Pubmed ID of the article

year Year of publication

journal Journal

Value

Object of class tbl_df, tbl, data.frame.

Source

Code for extracting p-values at: [inst/script/getPvalues.R](#)

| | |
|--------|---|
| lm_pi0 | <i>Estimation of π_0, proportion of p-values consistent with a null hypothesis</i> |
|--------|---|

Description

Estimation of π_0 , proportion of p-values consistent with a null hypothesis

Usage

```
lm_pi0(
  p,
  lambda = seq(0.05, 0.95, 0.05),
  X,
  type = c("logistic", "linear"),
  smooth.df = 3,
  threshold = TRUE,
  smoothing = c("unit.spline", "smooth.spline")
)
```

Arguments

| | |
|-----------|---|
| p | numeric vector, p-values |
| lambda | numeric vector, thresholds used to bin pvalues, must be in [0,1). |
| X | numeric matrix, covariates that might be related to p values (one test per row, one variable per column). |
| type | character, type of regression used to fit features to pvalues |
| smooth.df | integer, degrees of freedom when estimating $\pi_0(x)$ with a smoother. |
| threshold | logical, if TRUE, all estimates are thresholded into unit interval; if FALSE, all estimates are left as they are computed |
| smoothing | character, type of smoothing used to fit π_0 |

Value

object of class 'lm_pi0', which is a list with several components

| | |
|-----------------|---|
| call | matched function call |
| lambda | numeric vector of thresholds used in calculating π_0 .lambda |
| X.names | character vector of covariates used in modeling |
| π_0 .lambda | numeric matrix of estimated $\pi_0(x)$ for each value of lambda. The number of columns is the number of tests, the number of rows is the length of lambda. |
| π_0 | numerical vector of smoothed estimate of $\pi_0(x)$. The length is the number of rows in X. |
| π_0 .smooth | (only output with smoothing="smooth.spline") Matrix of fitted values from the smoother fit to the $\pi_0(x)$ estimates at each value of lambda (same number of rows and columns as π_0 .lambda) |

Examples

```
# define a covariate
X <- seq(-1,2,length=1000)
# set probability of being null
pi0 <- 1/4*X + 1/2
# generate null/alternative p-values
nullI <- rbinom(1000,prob=pi0,size=1)> 0
```

```
# vector of p-values
pValues <- rep(NA,1000)
pValues[nullI] <- runif(sum(nullI)) # from U(0,1)
pValues[!nullI] <- rbeta(sum(!nullI),1,2) # from Beta
pi0x <- lm_pi0(pValues, X=X)
```

lm_qvalue

*Estimation of qvalues conditioned on covariates***Description**

The recipe for turning pvalues into qvalues is adapted from package 'qvalue' and articles by Storey, Tibshirani, Taylor, Siegmund.

Usage

```
lm_qvalue(
  p,
  X,
  pfdR = FALSE,
  pi0 = NULL,
  smoothing = c("unit.spline", "smooth.spline"),
  ...
)
```

Arguments

| | |
|-----------|--|
| p | numeric vector of p-values |
| X | matrix of covariates (can be missing if pi0 is specified instead) |
| pfdR | logical, making estimates robust for small p-values and a small sample size |
| pi0 | list with pi0 estimates from lm_pi0. If this is not provided, pi0 is estimated using function lm_pi0. |
| smoothing | character, type of smoothing used to fit pi0. Note the default in this function is different than in lm_pi0. |
| ... | other parameters (passed on to lm_pi0 if pi0 is not provided) |

Value

object of class 'lm_qvalue', which is a list with several components

| | |
|---------|--|
| call | matched function call |
| pvalues | numeric vector of original p-values |
| qvalues | numeric vector of q-values |
| | other list elements transferred from pi0 |

Examples

```
# define a covariate
X <- rep(c(0, 1), each=1000)
# generate p-values, randomly for group 0 and with low values for group 1
pVal <- c(runif(1000), rbeta(1000, 0.2, 1))
# compute qvalues, using the covariate
qVal <- lm_qvalue(pVal, X=X)
```


Index

* datasets

BMI_GIANT_GWAS_sample, [2](#)
journals_pVals, [5](#)

BMI_GIANT_GWAS_sample, [2](#)

calculateSwfdr, [3](#)

get_number_decimals, [4](#)

journals_pVals, [5](#)

lm_pi0, [5](#)

lm_qvalue, [7](#)